

FIG. 1A




FIG. 1B

Fig. 2A





Fig. 2B



09882606.064501

Fig. 3A




Fig. 3B




Fig. 4A





Fig. 4B



09882605 "061501

Fig. 5A




Fig. 5B




Fig. 6: HIV-1 Protease protein surface showing an actual ligand as determined by x-ray co-crystal structure and showing the pocket of highest volume as calculated by the method of this application




Fig. 7: Heat Shock Protein 90 surface showing an actual ligand as determined by x-ray co-crystal structure and showing the pocket of highest volume as calculated by the method of this application




Fig. 8: Stromelysin protein surface showing an actual ligand as determined by x-ray co-crystal structure and showing the pocket of highest volume as calculated by the method of this application




Fig. 9: Dihydrofolate Reductase protein surface showing an actual ligand as determined by x-ray co-crystal structure and showing the pocket of highest volume as calculated by the method of this application





FIG. 10



09882606-061501

FIG. 11




FIG. 12





FIG. 13



09882606 .06.1501

FIG. 14:





FIG. 15



US 2004/0150110 A1

FIG. 16.




FIG. 17




FIG. 18





Figure 19:
Two plane slices shown for $N = 4$ (four points PNT distributed about a sphere)



09882606-061501
TEST90-9088860